

Northern Leopard Frog - *Lithobates pipiens*

Abundance: Unknown

Status: NSSU

NatureServe: G5 S3

Population Status: Once widely distributed, but rangewide declines have been documented. Many populations may be recovering. Extirpation is not eminent.

Limiting Factor: Habitat: declines in habitat quality have resulted in increased mortality. Similar or increased levels of chemical runoff (pesticides, herbicides, and household chemicals) likely to continue. However, No one factor can be attributed to species decline.

Comment: Was ranked NSS4 due to unknown status . The genus of the species was changed from *Rana*. Formerly *Rana pipiens*.

Introduction

The Northern Leopard Frog is historically one of the most common and widespread anurans in the United States. However, populations are known to be declining throughout its range. Northern Leopard Frogs may be found throughout Wyoming, but have experienced documented declines in the Greater Yellowstone Ecosystem and Laramie plains. Northern Leopard Frogs become active after water temperatures exceed 50 degrees Fahrenheit. The breeding season can extend from mid-March through July. Females typically deposit 600-7,500 eggs in a tight oval mass (Rorabaugh 2005). Depending on temperature, larval metamorphosis often occurs 3-6 months following egg deposition. In higher elevations, tadpoles may not metamorphose before winter. In Wyoming, Northern Leopard Frogs are not believed to overwinter as tadpoles (Baxter and Stone 1985). Northern Leopard Frogs actively forage among sedges, cattails, and tall grasses. Primary food items are invertebrates (beetles, flies, ants, worms, snails, etc.); however adult frogs will sometimes consume voles, small birds, snakes, and other amphibians (Baxter and Stone 1985). Resting near pond and lake margins, this species will quickly leap into the water if alarmed. Northern Leopard Frogs are known to winter in ponds, buried in mud. In many cases, they use a shallow pond for breeding and deep pools to hibernate.

Habitat

The Northern Leopard Frog can be found in or near permanent water in the plains, foothills, and montane zones (Smith and Keinath 2007). They can range to over 8,500 feet in elevation. Their preferred habitats are swampy cattail marshes on the plains and beaver ponds in the foothills and montane zones.

Problems

- h Habitat changes and other factors may be adversely affecting this species, but lack of data precludes identification of specific problems and development of management recommendations
- h Population status, distribution, habitat data, and disease status are lacking for this species.
- h Alteration of aquatic habitats needed for breeding may adversely affect populations.

Conservation Actions

- h A systematic study of this species should be conducted with respect to distribution, abundance, habitat associations, and disease status within Wyoming.
- h Continue efforts to educate landowners and the public about the importance of amphibians.
- h Develop management recommendations based on survey data.

Monitoring/Research

Conduct baseline surveys to gain better understanding of species distribution within the state. Monitoring should be conducted on known populations to ensure species viability.

Recent Developments

Amphibians have received increased attention within Wyoming. Incidental observations are encouraged to be reported to the herpetology program. Baseline surveys have been conducted in SW Wyoming to better understand herpetofaunal assemblages and distribution. Northern Leopard Frog distribution and habitat associations are being examined. Northern Leopard Frog monitoring has been conducted in the Powder River Watershed in conjunction with Coal Bed Methane discharge. Baseline surveys have been conducted for this species in Bighorn National Forest


References

Baxter, G.T. and M.D. Stone. 1985. Amphibians and Reptiles of Wyoming. Second Edition. Wyoming Game and Fish Department, Cheyenne. 137pp.

Rorabaugh J. C.. 2005. *Rana pipiens* Schreber, 1782 Northern Leopard Frog. Pages 570-577 in M.J. Lannoo (ed), Amphibian Declines: The Conservation Status of United States Species. University of California Press, Berkeley, CA.

Smith, B.E. and D.A. Keinath. 2007. Northern Leopard Frog (*Rana pipiens*): a technical conservation assessment. USDA Forest Service, Rocky Mountain Region.

Northern Leopard Frog (*Lithobates pipiens*)



SOURCE: Digital maps of ranges and predicted distributions for Wyoming Species of Greatest Conservation Need: April 2010. Wyoming Natural Diversity Database. University of Wyoming, Laramie, Wyoming. Note that brown indicates the predicted distribution of the species; heavy black lines indicate outermost boundaries of possible occurrence.